

PART I. SUBJECT INDEX

- Abramis brama** recommended for introduction into Canadian waters, Borodin, N. A., 368.
- Acipenser ruthenus** recommended for introduction into Canadian waters, Borodin, N. A., 368.
- Adirondack** lakes, chemical characteristics, Faigenbaum, Harold M., 189-196.
- Age of trout** in relation to time of spawning, Dinsmore (disc.), 358-359.
- Agersborg's** trout food, excellent results with, 155-162.
- Agriculture** and aquiculture, Agersborg, H. P. Kjerschow, 266-269.
- Alewife**: see *Pomolobus pseudoharengus*.
- Algae**, factor in some hatchery mortalities, M'Gonigle, R. H., 416-422.
- influence on fluctuation of dissolved oxygen in fishponds, Wiebe, A. H., 181-188.
- American Fisheries Society** constitution and by-laws, as amended to date, 472-474.
- membership, 475-494.
- Anax junius** larvae possible effective elimination by goldfish, Langlois (disc.), 138.
- Anchor** parasite (*Lernaea carassii*) infesting goldfish and carp, Tidd, Wilbur M., 176-180.
- Angling**, need for measure of catch, Clark, G. H., 49-53.
- Aquatic** insects in Waddell Creek riffles, their contribution by weight and numbers to available food supply, Needham, P. R., 245-246.
- plants, dominating species in lakes Brome and Manitou, Quebec, Spier, J. D., 464-468.
- plants in relation to oxygen supply and their bearing on fish life, Roach, Lee S., and Wickliff, E. L., 370-376.
- vegetable organisms furnished with nutrient salts by flooding land areas, Huntsman, A. G., 366-367.
- Aquiculture** and agriculture, Agersborg, H. P. Kjerschow, 266-269.
- scientific methods of control suggested, Huntsman, A. G., 364-367.
- Arizona**, effect of 1934 drought on fish life, James, M. C., 61.
- Arkansas**, effect of 1934 drought on fish life, James, M. C., 60.
- largemouth black bass rearing at Lonoke, Hogan, Joe, 127-131.
- Artificial spawning** of speckled trout, use of physiological saline, Werner, W. H. R., 346-349.
- Atlantic salmon**, cause of high mortality after spawning, Belding, David L., 219-222.
- spawning habits, Belding, David L., 211-216.
- spring vs. fall running, Belding, David L., and Kitson, J. Arthur, 225-230.
- taxonomy, Mottley, C. McC., 324.
- Backswimmers** and beetle larvae eliminated with cod-liver oil spray, Oliver, R., 136.
- Bacterial count** and chemical tests, significance in determining relative freshness of haddock, Griffiths, Francis P., and Stansby, Maurice E., 401-406.
- Bait dealers** should propagate their own minnows to obviate wholesale destruction of forage fish, Markus, Henry C., 93-96.
- Baker**, Davis (In Memoriam), 46.
- Barr**, Harry, inventor of only satisfactory fishway tested, Webster, B. O., 30.
- "Barren"** rainbow trout spawning later showed New Hampshire spawning season to extend from last November to middle of June, Agersborg, H. P. Kjerschow, 168.

- Bass** breeding in fertilized ponds, ecological study, Meehan, O. Lloyd, 151-154.
- largemouth black, rearing at Lonoke, Arkansas, Hogan, Joe, 127-131.
- new hatchery at South Otselec, New York, and its first year's operations, Kingsbury, Oliver R., 132-138.
- production costs cut materially by floating equipment, Regan, C. C., 143-145.
- production, standard methods of computing, Langlois, T. H., 163-166.
- social behavior in rearing ponds, Langlois, T. H., 146-150.
- Beetle** larvae and backswimmers eliminated with cod-liver oil spray, Oliver, R., 136.
- Belleville** hatchery, Bay of Quinte, maskinonge culture, MacKay, H. H. and Werner, W. H. R., 313-317.
- Black bass** legislation (1935), Denmead, Talbott, 97-102.
- Black-head minnows** (*Pimephales promelas*) should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- Black-nosed** dace, aquatic and surface drift food eaten, tabulated, 73.
- Black River** and Oswegatchie watershed, chemical characteristics, Faigenbaum, Harold M., 189-196.
- Blood suckers**, do they affect the fish population of a lake? Johnson, Herbert, 363.
- Blunt-nose minnows** (*Hyborhynchus notatus*) should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- Bottom samples**, method of taking, Needham, P. R., 238-240.
- Bream**, *Abramis brama*, recommended for introduction into Canadian waters, Borodin, N. A., 368.
- as affected by salt water in varying concentrations, survival in hours tabulated, 84.
- Breeding** of older males recommended, Agersborg, H. P. Kjerschow, 267.
- selective, for growth promotion in trout, Davis, H. S., 197-202.
- British Columbia** studies on coarse fish and the predator problem, Clemens, W. A., 318-321.
- Brome lake, Quebec**, dominating species of aquatic plants, Spier, J. D., 464-468.
- plankton distribution, Phillips, Mrs. J. T., 461-463.
- Brook trout** growth when released, vs. growth in hatchery (disc.), 80.
- spawning period, White, H. C., 356-357.
- aquatic and surface drift food eaten, in tabular form, 73.
- Buckeye Lake: see **Lake Buckeye**.
- Bullheads**, aquatic and surface drift food eaten, in tabular form, 74.
- By-laws** of American Fisheries Society as amended to date, 472-474.
- Canada**, introduction of Russian fishes recommended, Borodin, N. A., 368-369.
- Cannibalism** in bass rearing, prevention methods, Langlois, T. H., 146-150.
- elimination in largemouth black bass rearing ponds, Hogan, Joe, 128.
- methods of preventing, Langlois (disc.), 139.
- Carassius auratus** infested with *Lernaea carassii*, Tidd, Wilbur M., 176-180.
- carassius recommended for introduction into Canadian waters, Borodin, N. A., 368.
- Carp**: see **Carassius auratus** and **Carassius carassius**.
- Cary**, Guy (In Memoriam), 46.
- Casselman**, E. S. (In Memoriam), 46.
- Census** of angler's catch, present methods inadequate (disc.), 51-53.
- Champlain watershed**, chemical characteristics, Faigenbaum, Harold M., 189-196.

- Chemical tests** and bacterial count in determining relative freshness of haddock, Griffiths, Francis P., and Stansby, Maurice E., 401-406.
- Chlorine gas** effective in killing coarse fish (disc.), 280.
- Circular pools** advantageous for bass raising, Surber, Eugene (disc.), 141.
- Cisco**, origin and distribution in inter-glacial period, Mottley, C. McC. 324-325.
- Cladocera** and Copepoda, distribution in Manitou and Brome lakes, Quebec, Phillips, Mrs. J. T., 461-463.
- Clupea pallasii** spawning, quantitative studies, Hart, John Lawson, and Tester, Albert L., 307-312.
- Coarse fish** eliminated previous to stocking, Catt, James, 276-280.
and predator problem in relation to fish culture, Clemens, W. A., 318-321.
- Coastal fisheries** and weather conditions, forecasting feasible, Hachey, H. B., 382-389.
- Code**, N.R.A., for the fishery industry, Fiedler, R. H., 32-36.
- Colorado**, effect of 1934 drought on fish life, James, M. C., 60.
- Combat** of male salmon at spawning time (disc.), Greeley, 217-218.
- Commercial fisheries**, value of questionnaires, Van Oosten, John, 107-117.
fishing, report division of, Clemens, W. A. (including Fiedler statement), 32-37.
- Committee** on American Fish Policy, report for 1933-1934, 25-27.
on Foreign Relations, report for 1933-1934, 20-23.
on relations with Federal, Provincial, and State governments, report for 1933-1934, 23-25.
- Conservation** of water, Bell, Frank T., 54-56.
- Constitution** and by-laws of American Fisheries Society, as amended to date, 472-474.
- Copepoda** and Cladocera, distribution in Manitou and Brome lakes, Quebec, Phillips, Mrs. J. T., 461-463.
- Copper sulphate** in elimination of coarse fish, Catt, James, 276-280.
- Coregonus clupeaformis** growth in Trout Lake, Wisconsin, Hile, Ralph, and Deason, Hilary J., 231-237.
albula, recommended for introduction into Canadian waters, Borodin, N. A., 368.
- Corliss**, C. G. (In Memoriam), 46.
- Cottonwood** lakes, California, life-history of golden trout, Curtis, Brian, 259-265.
- Counting fry** and fingerling rapidly and accurately, new method, Prevost, Gustave, 270-275.
- Crayfish** in bass ponds for control of vegetation (disc.), 150.
- Creek chub** spawning habits; its propagation in ponds, Markus, Henry C., 95.
- Cut-throat**, taxonomy, Mottley, C. McC., 324.
- Cyprinus carpio** infested with Lernaecarassii, Tidd, Wilbur M., 176-180.
- Dace**: see **horned dace**, **black-nosed dace**.
- Dams** on lakes and streams, their physiographical and mechanical effects on the impounded waters, Richardson, L. R., 457-458.
- Daphnia** cultures at new South Otselic hatchery, New York, Kingsbury, Oliver R., 135-136.
and other forage organisms propagated intensively in small ponds, Embury, G. C., and Sadler, W. O., 205-210.
pulex survived in water with an oxygen tension down to 0.05 c.c. per litre, Smith, M. W., 414.
- Deforestation**, ill effects in Mississippi River Valley, Culler, C. F., 328.
- Departure Bay**, herring spawning, quantitative studies, Hart, John Lawson, and Tester, Albert L., 307-312.
- Disease** of fish, ulcer, described, Fish, Frederic F., 252-258.

- fungous, appearance on eggs coincident with desertion of the nest by the male, Kingsbury, Oliver R., 137.
- Diseases** of fish, treatment of underlying causes rather than symptoms to be desired, Huntsman, A. G., 1934, 364-367.
- Distribution** of bottom foods, factors governing, Needham, P. R., 246-247.
- Dolly Varden**, origin and distribution in inter-glacial period, Mottley, C. McC., 324-325.
- Domestication** of bass to prevent cannibalism, Langlois, T. H., 146-150.
- Domesticated** vs. wild trout, differing rates of growth, Dinsmore, A. H., 203-204.
- Drought** of 1934, effect on fish life, James, M. C., 57-62.
- Dunlap**, Irving H. (In Memoriam), 46.
- Eaton**, E. H. (In Memoriam), 46.
- Ecology** of fertilized ponds for bass rearing, Meehan, O. Lloyd, 151-154.
- Egg planting**, "green" vs. eyed-eggs, Foerster, R. E., 381.
- Eggs as food**, best method of feeding, Agersborg, H. P. Kjerschow, 440.
of herring at Departure and Nanoose Bays, Vancouver Island, number spawned and mortality, Hart, John Lawson, and Tester, Albert L., 307-312.
of trout, relation of temperature to incubation period, Embody, G. C., 281-292.
- European** trout, origin and taxonomy, Mottley, C. McC., 324-325.
- Environment** vs. heredity as influence on time of spawning, Greeley (disc.), 358.
- Environmental conditions** in trout stream management, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 68-80.
- Evolution** in salmonoids, Mottley, C. McC., 323-327.
- Feeding habits** of maskinonge, MacKay, H. H., and Werner, W. H. R., 315-316.
of males for more abrupt and complete ripeness, Agersborg, H. P. K., 267.
- Fertilization** of ripe ova with milt from salmon parr produced fry with 60 per cent deformity and eyeless monsters, White, H. C., 361.
- Fertilized** waters, dissolved oxygen content, Smith, N. W., 408-415.
- Fertilizers**, dried sludge from disposal plant successfully used in equal amounts with cow manure, Markus (disc.), 436.
influence on dissolved oxygen in fish ponds, Wiebe, A. H., 181-188.
in pondfish production, ecological aspects, Meehan, O. Lloyd, 151-154.
in ponds, early vs. late fertilizing, Hogan (disc.), 140.
in rearing ponds for largemouth black bass, Hogan, Joe, 129.
used in Daphnia and other forage organism culture, Embody, G. C., and Sadler, W. O., 205-210.
used at new South Otselic hatchery, New York, Kingsbury, Oliver R., 134-136.
- Fin rot disease** and ulcer, differences shown, Fish, Frederic F., 255.
- Fingerling** and fry counting, new method, Prevost, Gustave, 270-275.
- Fish culture**, better staff recommended to obviate present poor success in aquaculture due to ignorance, Agersborg, H. P. Kjerschow, 268-269.
report Vice-President Division of, for 1933-1934, Webster, B. O., 28-32.
- Fish** in fresh water streams as affected by salt water from oil wells, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 81-86.
- Fish life** as affected by drought of 1934, James, M. C., 57-62.
- Fish meal**, white, nutritive value, comparison of laboratory and practical tests for determining, Cleveland, M. M., and Fellers, C. R., 293-303.
- Fish species** suggested for rearing in sewage treatment ponds, Radebaugh, Gus H., and Agersborg, H. P. Kjerschow, 454.

- Fisheries** in coastal regions and weather forecasting, Hachey, H. B., 382-389.
- Fishways**, discussion of various types, including the new Barr fishway, Webster, B. O., 29-30.
- Flooding** areas, mechanical as well as chemical aspects to be considered, Richardson (disc.), 367.
- Food** in bottom samples, relation to stream width, variability, seasonal distribution, type vs. quantity, Needham, P. R., 238-240.
- cost of trout in New York State hatcheries, Deuel, Charles R., 172-175.
- eaten by trout liberated from hatcheries, tabulated, Lord, Russell F., 339-345.
- nutritive value of ground fish meals, Cleveland, M. M., and Fellers, C. R., 293-303.
- Food for fish:** beef-heart, beef-liver and Swift's beef meal followed by beef liver and salmon egg meal for young bass, Surber, Eugene (disc.), 141.
- comparison of diets made at South Otselic bass hatchery, Kingsbury, Oliver R., 136.
- fresh, vs. canned carp in bass hatcheries, Langlois (disc.), 139.
- increased production in stream improvement, Davis H. S., 63-66.
- inexpensive balanced diet for trout and salmon, Agersborg, H. P. K., 155-162.
- quantitative and qualitative analyses of fresh commercial fisheries products for artificial rearing of salmonoids, Agersborg, H. P. K., 435-442.
- summary of aquatic and surface drift food, tabulated, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 73-74.
- Food grinder**, new type to prevent pulping of feed, Langlois (disc), 162
- Forage fish**, wholesale destruction, Markus, Henry C., 93-96.
- organisms propagated intensively in small ponds, Embody, G. C., and Sadler, W. O., 205-210.
- Foraging** of liberated trout, stomach contents analyzed, Lord, Russell F., 339-345.
- Foreign** Relations Committee, report for 1933-1934, Rodd, J. A., 20-23.
- Freshwater fish**, effect of salt water environment, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 81-86.
- Fry** and fingerling counting, new method, Prevost, Gustave, 270-275.
- production from eyed-egg planting, Foerster, R. E., 379-381.
- Fungous** growth on eggs coincident with desertion of the nest by the male, Kingsbury, Oliver R., 137.
- Furunculosis**, different from ulcer disease, Fish, Frederic F., 252.
- Gaspé County, Quebec**, general discussion of lakes, Taylor, B. W., and Lindsay, R. C., 424-431.
- Glenora hatchery**, Bay of Quinte, observations on culture of maskinonge, MacKay, H. H., and Werner, W. H. R., 313-317.
- Golden shiner** as affected by salt water in varying concentrations, survival in hours tabulated, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 84.
- should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- Golden trout** of Cottonwood lakes, life-history, Curtis, Brian, 259-265.
- Goldfish** infested by *Lernaea carassii*, Tidd, Wilbur M., 176-180.
- as possible effective means of eliminating dragonfly larvae, Langlois (disc.), 138.
- Grayling**, origin and distribution in inter-glacial period, Mottley, C. McC., 324-325.

- Ground fish meals**, nutrient value, comparison of laboratory and practical tests for determining, Cleveland, M. M., and Fellers, C. R., 293-303.
- Growth** of brook trout, effect of heredity, Dinsmore, A. H., 203-204.
- of brook trout in hatcheries, vs. growth when released, Markus, H. C. (disc.), 80.
- of chicks, red fish meal as sole source of protein, experiments, Cleveland, M. M., and Fellers, C. R., 297-301.
- of golden trout in Cottonwood lakes, California, Curtis, Brian, 259-265.
- and heredity in trout, Davis, H. S., 197-202.
- of *Pomolobus pseudoharengus* in Seneca Lake, N. Y., Odell, T. T., 118-126.
- rate of trout as studied in Trammel Creek, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 74-76.
- of rats fed fish meal as sole source of protein, experiments, Cleveland, M. M., and Fellers, C. R., 295-297.
- of salmon parr, improved technical methods for determining by scale measurements, Belding, David L., 103-106.
- of selected trout in Antigonish hatchery, Nova Scotia (disc.), 202.
- of trout treated with potassium permanganate vs. treated with salt, Prevost, Gustave, 304-306.
- and water requirements of maskinonge, MacKay, H. H., and Werner, W. H. R., 316-317.
- of whitefish in Trout Lake, Wisconsin, Hile, Ralph, and Deason, Hilary J., 231-237.
- Haddock** freshness, significance of bacterial count and chemical tests in determination, Griffiths, Francis P., and Stansby, Maurice E., 401-406.
- Hatchery** for bass, floating, Regan, C. C., 143-145.
- for bass at South Otselic, N. Y., and its first year's operations, Kingsbury, Oliver R., 132-138.
- predators, Hogan, Joe, 129.
- fry emerge during day-light, whereas experiment with planted eye-eggs in Cultus Lake showed the fry to emerge from the gravel bed between dusk and dawn, Foerster, R. E., 390.
- shown to have little evil effect on the gallant nature of trout, experiments described, Lord, Russell F., 339-345.
- Haynes**, W. deF. (In Memoriam), 46.
- Health** coefficient much greater with new diet than with beef liver or any other trout food used, Agersborg, H. P. K., 160.
- Heredity**, effect on growth of brook trout, Dinsmore, A. H., 203-204.
- and growth in trout, Davis, H. S., 197-202.
- vs. environment as influence on time of spawning, Greeley (disc.), 358.
- Herring meal** fertilizer effect on oxygen content in open containers shown graphically, Smith, M. W., 411.
- spawning, quantitative studies, Hart, John Lawson, and Tester, Albert L., 307-312.
- Hogarth**, George R. (In Memoriam), 46.
- Horned dace** (*Semotilus atromaculatus*) should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- summary of aquatic and surface drift food eaten, tabulated, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 73.
- Hyborhynchus** notatus should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- nuchalis, wholesale destruction, Markus, Henry C., 93-96.
- Hydrogen-ion concentration** as influenced by algae, M'Gonigle, R. H., 420.

- in ponds as influenced by vegetation and fertilizers, Wiebe, A. H., 181-188.
- Idaho**, effect of 1934 drought on fish life, James, M. C., 61.
- Illinois**, effect of 1934 drought on fish life, James, M. C., 59.
- Immature salmon parr** fertilize ripe ova, but 60 per cent of resulting fry were deformed or eyeless monsters, White, H. C., 361.
- Impounded waters**, effects of dams on lakes and streams, ecology, physiographical and mechanical elements involved, Richardson, L. R., 457-458.
- in water conservation, Bell, Frank T., 55.
- Impounding** of salt waters from oil fields in Texas, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 81-86.
- In **Memoriam**, 46.
- Incubation** period of trout eggs, effect of temperature, Embury, G. C., 281-292.
- Indiana**, effect of 1934 drought on fish life, James, M. C., 58.
- Insects**, aquatic, in Waddell Creek riffles, their contribution by weight and numbers to available food supply, Needham, P. R., 245-246.
- Introduction** into Canadian waters of Russian fish recommended, Borodin, N. A., 368-369.
- Iowa**, effect of 1934 drought on fish life, James, M. C., 59.
- nursery pond kettle and outlet, Aitken, W. W., 170-171.
- Izaak Walton League**: improvement of pollution control, Locke, S. B., 87-92.
- Kamloops** and steelhead physiographical lines of separation, Mottley, C. McC., 326-327.
- Kansas**, effect of 1934 drought on fish life, James, M. C., 61.
- Kettle and elbow outlet, new type in Iowa nursery pond, Aitken, W. W., 170-171.
- Laboratory** for experimental research on effect of salt water on fish, organized in Texas, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 83.
- vs. practical tests for determining the nutritive value of fish meals, Cleveland, M. M., and Fellers, C. R., 293-303.
- Lake** bottom improvement, discussion in report Vice-President Division of Fish Culture, Webster, B. O., 31.
- Buckeye, Ohio, aquatic vegetation and its bearing on fish life, Roach, Lee S., and Wickliff, E. L., 370-376.
- Brome, Quebec, dominating species of aquatic plants, Spier, J. D., 464-468.
- Brome, Quebec, plankton distribution, Phillips, Mrs. J. T., 461-463.
- Champlain, chemical characteristics, Faigenbaum, Harold M., 189-196.
- Cultus experiments in planting 50,000 eyed-eggs with resultant production of 44,691 fry (88% of total eggs planted), Foerster, R. E., 379-381.
- Huron, average spawning seasons for lake trout, whitefish, yellow pike-perch and perch, tabulated, Van Oosten, John, 112.
- Manitou, Quebec, dominating species of aquatic plants, Spier, J. D., 464-468.
- Manitou, Quebec, plankton distribution, Phillips, Mrs. J. T., 461-463.
- Michigan, average spawning seasons for lake trout, whitefish, yellow pike-perch and perch, tabulated, Van Oosten, John, 112.
- Lake Superior**, average spawning seasons for lake trout, whitefish, yellow pike-perch, and perch, tabulated, Van Oosten, John, 112.
- Wapello rearing ponds, new type kettle and outlet, Aitken, W. W., 170-171.
- Lake trout**, origin and distribution in inter-glacial period, Mottley, C. McC., 324-325.
- spawning season, in lakes Huron, Michigan, and Superior, tabulated, Van Oosten, John, 112.

- Lakes of Gaspe County, Quebec**, general discussion, Taylor, B. W., and Lindsay, R. C., 424-431.
- 200-300 acres to be stocked, when not fed by considerable waters containing enemy fish, are treated with copper sulphate to eliminate coarse fish, Catt, James, 276-280.
- Largemouth black bass**, 1935 legal requirements for further protection, Denmead, Talbott, 97-102.
- rearing at Lonoke, Arkansas, Hogan, Joe, 127-131.
- Laws and regulations: see **Legislation**.
- Legislation** affecting fishery industry enacted during the 73rd Congress, 2nd Session, Clemens, W. A., in report Division of Commercial Fishing for 1933-1934, 37.
- for black bass (1935), Denmead, Talbott, 97-102.
- Lernaea carassii** ("anchor parasite") infesting goldfish and carp, Tidd, Wilbur M., 176-180.
- Liberating** young salmon in a stream where salmon had never run brought spawning salmon the following spring, White, H. C., 361.
- Librarian's report**, Cobb, Eben W., 19.
- Ling**, origin and distribution in inter-glacial period, Mottley, C. McC., 324-325.
- Manitou lake, Quebec**, dominating species of aquatic plants, Spier, J. D., 464-468.
- plankton distribution, Phillips, Mrs. J. T., 461-463.
- Maskinonge** culture at Belleville and Glenora fish hatcheries, Bay of Quinte, MacKay, H. H., and Werner, W. H. R., 313-317.
- Mating**, constant discarding of older males for younger specimens detrimental, Agersborg, H. P. Kjerschow, 267.
- of salmon described, Belding, David L., 213-214.
- Members** American Fisheries Society, honorary, 475; patrons, 475-476; life, 477; active, 478-492; state membership, 492; clubs, dealers, etc., 492-493; libraries, 494; corresponding, 494.
- Michigan**, effect of 1934 drought on fish life, James, M. C., 57.
- recommendations on regulations based on majority opinion of commercial fishermen, Van Oosten, John, 115.
- spawning season for lake trout, whitefish, yellow pike-perch, tabulated, Van Oosten, John, 112.
- Migration** of salmon, factors influencing their return from the sea, Huntsman, A. G., 351-354.
- of steelhead and silver salmon, Taft, A. C., 251.
- Milt** from immature salmon parr fertilized eggs, but 60% of resulting fry were deformed or eyeless monsters, White, H. C., 361.
- Minnesota**, installation of sewage disposal plants at Minneapolis and St. Paul, 28-29.
- Minnows** should be reared by bait dealers to obviate their wholesale destruction, Markus, Henry C., 93-96.
- Mississippi** fisheries, deforestation, erosion, etc., recommended corrections, Culler, C. F., 328-329.
- Missouri**, effect of 1934 drought on fish life, James, M. C., 57.
- Mortality** of Atlantic salmon after spawning, cause, Belding, David L., 219-222.
- in trout and salmon reduced to practically zero, Agersborg, H. P. Kjerschow, 160.
- Muskellunge** propagation in Wisconsin, Webster, B. O., 31-32.
- N. R. A.** code for fish industry, Fiedler, R. H., 32-36.
- NanOOSE Bay** herring spawning, quantitative studies, Hart, John Lawson, and Tester, Albert L., 307-312.
- Nebraska**, effect of 1934 drought on fish life, James, M. C., 59.
- Nests**, floating, for bass, Regan, C. C., 143-145.

- New York, new bass hatchery at South Otselic, and its first year's operations, Kingsbury, Oliver R., 132-138.
- state hatcheries, cost of trout food, Deuel, Charles R., 172-175.
- Nominations of officers, 43.
- Notemigonus crysoleucas* should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- Nutritive value of white fish meals, Cleveland, M. M., and Fellers, C. R., 293-303.
- Oceanography meteorology and the fisheries, Hachey, H. B., 382-389.
- Ohio, effect of 1934 drought on fish life, James, M. C., 58.
- Oklahoma, effect of 1934 drought on fish life, James, M. C., 59.
- Omerus mordax* Mitchill, method of planting on a small scale, Richardson, L. R., and Belknap, G. W., 432-434.
- Oswegatchie and Black River watershed, discussion of depth, temperature, free carbon dioxide and methyl orange alkalinity, dissolved oxygen and pH of bottom waters, Faigenbaum, Harold M., 189-196.
- Outlet, new type in Iowa nursery pond, Aitken, W. W., 170-171.
- Oxygen content of artificial lakes with a great deal of timber (disc.), 459-460.
- in fertilized waters, Smith, M. W., 408-415.
- of water ranging from .15 to 2.00 parts per million, actual case of fish surviving for several hours, Wickliff (disc.), 376.
- Oxygen depletion by algae cause of fish mortality in hatcheries, M'Gonigle, R. H., 416-422.
- and vegetation control in shallow waters, Hayford, Charles O. (disc.), 377.
- Oxygen, nocturnal depressions in fish ponds with special reference to excess vegetation and fertilizers, Wiebe, A. H., 181-188.
- supply of aquatic plants and their bearing on fish life, Roach, Lee S., and Wickliff, E. L., 370-376.
- Parasites and their importance, Parnell, I. W., 390-399.
- Parasitism: see Blood suckers.
- Parr: see *Salmon parr*.
- Perch, spawning season in lakes Huron, Michigan, and Superior, tabulated, Van Oosten, John, 112.
- Perrier, Edmond (In Memoriam), 46.
- Physiological saline in experiments with artificial spawning of speckled trout, Werner, W. H. R., 346-349.
- Phytoplanktonic growth, effect on oxygen content in fertilized water shown graphically, Smith, M. W., 412.
- Pike, origin and distribution in inter-glacial period, Mottley, C. McC., 324-325.
- Pike-perch, yellow, spawning season in lakes Huron, Michigan and Superior, tabulated, Van Oosten, John, 112.
- Pimephales *promelas* should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- Plankton distribution in Manitou and Brome lakes, Quebec, Phillips, Mrs. J. T., 461-463.
- Planting of *Osmerus mordax* Mitchill on a small scale, method described, Richardson, L. R., and Belknap, G. W., 432-434.
- Pollution control in Michigan (disc.), 91-92.
- federal agency for control proposed as against present inadequate state provisions, Locke, S. B., 87-91.
- of hatchery waters by algae causing severe annual loss of fish, M'Gonigle, R. H., 416-422.
- of streams by salt water from oil fields, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 81-86.
- Pomolobus pseudoharengus*, life history and ecology in Seneca Lake, New York, Odell, T. T., 118-126.

- Potassium permanganate** in fish culture, a criticism, Prevost, Gustave, 304-306.
- Predator** and coarse fish problem in relation to fish culture, Clemens, W. A., 318-321.
- Predators** (grebe, herons, terns, kingfishers and snakes) at Lonoke hatchery, Arkansas, Hogan, Joe, 129.
- Propagating** daphnia and other forage organisms intensively in small ponds, Embury, G. C., and Sadler, W. O., 205-210.
- Propagation**, stream improvement in relation to, Davis, H. S., 63-67.
- Protection** of black bass, 1935 legal requirements, Denmead, Talbott, 97-102. and legislation, report division of, Amsler, Guy, 38-39.
- Proteus hydrophilus** possibly secondary invader in ulcer disease lesions, Fish, Frederic F., 253.
- Quebec**, Lakes Manitou and Brome, plankton distribution, Phillips, Mrs. J. T., 461-463; Spier, J. D., 464-468.
- trout lakes in Gaspé County, general discussion, Taylor, B. W., and Lindsay, R. C., 424-431.
- Rainbow trout**, origin, taxonomy and relations, Mottley, C. McC., 323-327.
- Raquette watershed**, chemical characteristics, Faigenbaum, Harold M., 189-196.
- Ravenel**, W. de C. (In Memoriam), 46.
- Reclamation**, hydroelectric, Bell, Frank T., 54-56.
- Redd** digging of salmon described, Belding, David L., 212-213.
- Regulations**, recommendations based on majority opinion of the commercial fishermen, State of Michigan, Van Oosten, John, 115.
- Rock bass**, environment vs. temperature as influence on time of spawning, Langlois (disc.), 359.
- Russian fishes** recommended to be introduced into Canadian waters, Borodin, N. A., 368-369.
- Saint Lawrence watershed**, chemical characteristics, Faigenbaum, Harold M., 189-196.
- Salinity**, low, not a factor influencing salmon toward river mouths, Huntsman (disc.), 355.
- Salmo** aqua-bonita Jordan, life-history; S. roosevelti, a color variation or subspecies of S. aqua-bonita, Curtis, Brian, 259-265.
- fontinalis spawning period, White, H. C., 356-357.
- Salmon** attracted to streams by milt in water? White, H. C., 360-361.
- factors influencing their return from the sea, Huntsman, A. G. 351-354.
- inexpensive balanced diet for, Agersborg, H. P., Kjerschow, 155-162.
- parr annual growth, improved technical methods for determining by scale measurements, Belding, David L., 103-106.
- regulations adopted for inspection and grading of canned, Clemens, W. A., in report Division of Commercial Fishing, 36.
- run on the Skagit after experimental planting of sockeye, Dinsmore (disc.), 362.
- secondary sexual characters in (disc.), 217.
- spawning habits, Belding, David L., 211-216.
- taxonomy, Mottley, C. McC., 323-324.
- varieties: evidence pro and con separate races, Belding, David L., and Kitson, J. Arthur, 225-230.
- Salt** treatment vs. potassium permanganate as influencing growth of trout, Prevost, Gustave, 304-306.

- water from oil fields polluting streams in Texas, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 81-86.
- Scale measurements**, improved technical methods in determination of annual growth of salmon parr, Belding, David L., 103-106.
- Scott Creek**, California, steelhead experiments, Taft, A. C., 248-251.
- Schuil**, Henry A. (In Memoriam), 46.
- Sea** conditions and atmospheric pressure,—weather and fishery forecasting, Hachey, H. B., 382-389.
- Secondary sexual characters** in salmon, Belding, David L. (disc.), 217.
- Selective breeding** in trout for growth promotion, Davis H. S., 197-202. of speckled trout, Shillington, K. G., 274-275.
- Semotilus atromaculatus** should be reared by bait dealers to obviate wholesale destruction of forage fish, Markus, Henry C., 94.
- Seneca Lake**, New York, ecological conditions with special reference to *Pomolobus pseudoharengus*, Odell, T. T., 118-126.
- Sewage treatment**, economic value of effluent in wildlife conservation, Radebaugh, Gus H., and Agersborg, H. P. Kjerschow, 443-445.
- Sheldon**, Edward D. (In Memoriam), 46.
- Shelter provision**, in stream improvement, Davis, H. S., 63-64.
- Shrimp**, exclusive diet of ground shrimp at South Otselic bass hatchery resulted in weakened and parasitized fish which quickly recovered with change of diet, Kingsbury, Oliver R., 136.
- Silt erosion** control essential factor, Culler, C. F., 328-329.
- Silver hake**, adult, an exclusive diet of one and one half inch brook trout, found detrimental, Agersborg, H. P. K., 155.
- Smallmouth black bass**, legal requirements for further protection, Denmead, Talbott, 97-102.
- Snake**: see **Water snake**.
- Soil erosion** control, Davis, H. S., 63-65.
- South Dakota**, effect of 1934 drought on fish life, James, M. C., 58.
- South Otselic**, New York, new bass hatchery and its first year's operations, Kingsbury, Oliver R., 132-138.
- Spawning**, artificial, comparison of fertility percentages using "dry" vs. physiological or normal saline solution methods, Dinsmore (disc.), 349.
- artificial, use of physiological saline in experiments, Werner, W. H. R., 346-349.
- of Atlantic salmon, first description of natural spawning, Belding, David L., 214-216.
- habits of steelhead trout, Needham, P. R., and Taft, A. C., 332-338.
- of herring, quantitative studies, Hart, John Lawson, and Tester, Albert L., 307-312.
- loss of weight previous to and during in Atlantic salmon, Belding, David L., 219-222.
- migration of steelhead and silver salmon, Taft, A. C., 251.
- of *Pomolobus pseudoharengus* in Seneca Lake, New York, Odell, T. T., 118.
- season in Michigan, in tabular form, from questionnaires, Van Oosten, John, 112.
- season of rainbow trout, Agersborg, H. P. K., 167-169.
- of trout, age as factor influencing period, Dinsmore (disc.), 358-359.

- of wild bass at new South Otselic hatchery, Kingsbury, Oliver R., 132.
- Speckled trout** selective breeding, Shillington, K. G., 274-275.
- Spermatozoa** of speckled trout kept alive for four minutes in normal saline, Werner, W. H. R., 346-349; activity, Greeley (disc.), 350.
- Sport** and alertness in trout liberated from hatcheries, Lord, Russell F., 339-345.
- State action** vs. federal agency to control pollution, Locke, S. B., 90.
- Steelhead** and silver salmon migration, Taft, A. C., 248-251.
spawning habits, Needham, P. R., and Taft, A. C., 332-338.
taxonomy, Mottley, C. McC., 323.
- Stewart**, William D. (In Memoriam), 46.
- Stream** bottom foods, Needham, P. R., 238-247.
improvement, discussion in report Vice-President Division of Fish Culture, Webster, B. O., 30-31.
improvement, purpose and value, Davis, H. S., 63-67.
management as studied in Trammel Creek, New York, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 68-80.
pollution with salt water from oil fields in Texas, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 81-86.
- Stunting** vs. growth inhibition in starved fish, Agersborg, H. P. Kjerschow, 439-440.
- Sturgeon**, *Acipenser ruthenus*, recommended for introduction into Canadian waters, Borodin, N. A., 368.
- Squawfish**, distribution in inter-glacial period, Mottley, C. McC., 325.
- Surveys**, value of questionnaires, Van Oosten, John, 107-117.
- Temperature** effect on incubation period of trout eggs, Embody, G. C., 281-292.
for rearing bass (disc.), 142.
requirements of *Pomolobus pseudoharengus*, Odell, T. T., 124.
of sea, factor influencing maturing of salmon, Huntsman, A. G., 351-354.
of spawning ponds for largemouth black bass, Hogan, Joe, 127.
of streams, control of, Davis, H. S., 63-66.
- Tench**, *Tinca tinca*, recommended for introduction into Canadian waters, Borodin, N. A., 368.
- Texas** stream pollution with salt water from oil fields, Wiebe, A. H., Burr, J. G., and Faubion, H. E., 81-86.
- Timber**, lakes containing quantities, and its effect on the oxygen content (disc.), 459-460.
- Tinca tinca** recommended for introduction into Canadian waters, Borodin, N. A., 368.
- Toxic** condition of water due to growth of algae (disc.), 423.
- Trammel Creek** (Mohawk watershed), New York, study of stream and suggested methods of trout stream management, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 68-80.
- Treasurer's report** for 1933-1934, 15-18.
- Trematode worms**, salt vs. potassium permanganate used in treatment for, influence on growth of trout, Prevost, Gustave, 304-306.
- Trexler**, Harry C. (In Memoriam), 46.
- Trout**, age in relation to time of spawning, Dinsmore (disc.), 358-359.

- brook, effect of heredity on growth, Dinsmore, A. H., 203-204.
brook, spawning period, White, H. C., 356-357.
eggs, relation of temperature to incubation period, Embody, G. C., 281-292.
feeding in New York State hatcheries, cost of, Deuel, Charles R., 172-175.
golden, in Cottonwood lakes, California, life-history, Curtis, Brian, 259-265.
growth and heredity, Davis, H. S., 197-202.
growth rate as studied in Trammel Creek, New York, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 74-76.
from hatcheries as foragers and game fish, Lord, Russell F., 339-345.
lakes in Gaspé County, Quebec, general discussion, Taylor, B. W., and Lindsay, R. C., 424-431.
Lake, Wisconsin, growth of whitefish (*Coregonus clupeaformis*), Hile, Ralph, and Deason, Hilary J., 231-237.
rainbow, origin and relations, Mottley, C. McC., 323-327.
rainbow, spawning season, Agersborg, H. P. Kjerschow, 167-169.
reared to legal size in less than ten months on new diet, Agersborg, H. P. K., 155-162.
speckled, selective breeding, Shillington, K. G., 274-275.
speckled, spermatozoa kept alive four minutes in normal saline in artificial spawning experiments, Werner, W. H. R., 346-349.
steelhead, spawning habits, Needham, P. R., and Taft, A. C., 332-338.
stream management, Moore, Emmeline, Greeley, J. R., Greene, C. W., Faigenbaum, H. M., Nevin, F. R., and Townes, H. K., 68-80.
Ulcer disease of trout, Fish, Frederic F., 252-258.
Upper Hudson watershed, chemical characteristics, Faigenbaum, Harold M., 189-196.
Urbana Champaign sewage treatment works, lagoon used as fish pond, microscopic biota tabulated, Radebaugh, Gus H., and Agersborg, H. P. Kjerschow, 447.
Utah, effect of 1934 drought on fish life, James, M. C., 60.
Vegetation areas alternately flooded and allowed to revert to land conditions for production of nutrient salts through decomposition of land plants recommended, Huntsman, A. G., 366-367.
of Buckeye Lake, oxygen supply and its effect on fish life, Roach, Lee S., and Wickliff, E. L., 370-376.
control, Langlois (disc.), 139-140.
control to prevent cannibalism in bass, Langlois, T. H., 148-149.
influence on dissolved oxygen in fish ponds, Wiebe, A. H., 181-188.
of lakes Brome and Manitou, Quebec, dominating species, Spier, J. D., 464-468.
Waddell Creek, California, steelhead experiments, Taft, A. C., 248-251.
Ward, Robertson S. (In Memoriam), 46.
Water conservation, Bell, Frank T., 54-56.
fluctuation control in stream improvement, Davis, H. S., 63-65.
pH as influenced by algae, M'Gonigle, R. H., 420.
pollution, installation of sewage disposal plants, in report Vice-President Division of Fish Culture, 28-32.
snakes found to feed abundantly on tadpoles with slim traces of fish, Langlois (disc.), 140.

- Weather** conditions and coastal fisheries, forecasting feasible, Hachey, H. B., 382-389.
- Weaver**, Judd S. (In Memoriam), 46.
- Weed cutters** (disc.), 377-378.
- Weight**, loss of, in Atlantic salmon previous to and during spawning period, Belding, David L., 219-222.
- White**, E. Hamilton (In Memoriam), 46.
- Whitefish** (*Coregonus clupeaformis*), growth in Trout Lake, Wisconsin, Hile, Ralph, and Deason, Hilary J., 231-237.
origin and distribution in inter-glacial period, Mottley, C. McC., 324-325.
spawning season, in lakes Huron, Michigan and Superior, in tabular form, Van Oosten, John, 112.
- Wild** vs. domesticated trout, differing rates of growth, Dinsmore, A. H., 203-204.
- Wisconsin**, effect of 1934 drought on fish life, James, M. C., 57.
Trout Lake, growth of whitefish (*Coregonus clupeaformis*), Hile, Ralph, and Deason, Hilary J., 231-237.
- Wyoming**, effect of 1934 drought on fish life, James, M. C., 58.
- Yellow pike-perch** spawning season, in lakes Huron, Michigan and Superior, in tabular form, Van Oosten, John, 112.
- Zalsman**, P. G. (In Memoriam), 46.

PART II. AUTHOR INDEX

- Adams, M. P.:** Pollution of Michigan streams with brine from oil fields (disc.), 85-86.
Pollution and water conservation in Michigan (disc.), 91-92.
- Agersborg, H. P. Kjerschow:** Aquiculture and agriculture, 266-269.
Inexpensive balanced diet for trout and salmon, 155-162.
Quantitative and qualitative analyses of foods for artificially reared salmonoids, 435-442.
When do rainbow trout spawn? 167-169.
- Agersborg, H. P. K.,** joint author: see **Radebaugh, Gus H.**
- Aitken, W. W.:** Iowa nursery pond kettle and outlet, 170-171.
- Amsler, Guy:** Report Division Protection and Legislation for 1933-1934, 38-39.
- Bekling, David L.:** High mortality in Atlantic salmon after spawning, 219-222.
Salmon parr annual growth, improved technical methods for determining scale measurements, 103-106.
Spawning habits of Atlantic salmon, 211-216.
- Belding, David L.,** and **Kitson, J. Arthur:** Spring-run and fall-run Atlantic salmon, 225-230.
- Belknap, G. W.,** joint author: see **Richardson, L. R.**
- Bell, Frank T.:** Report Committee on Relations with Federal, Provincial, and State governments for 1933-1934, 23-25.
Water conservation, 54-56.
- Borodin, N. A.:** Introduction of Russian fishes into Canadian waters, 368-369.
- Burr, J. C.,** joint author: see **Wiebe, A. H.**
- Catt, James:** Copper sulphate in elimination of coarse fish, 276-280.
Growth of selected trout in Antigonish hatchery, Nova Scotia (disc.), 202.
- Clark, G. H.:** Need for a measure of the angler's catch, 49-53.
- Clemens, W. A.:** Predator and coarse fish problem in relation to fish culture, 318-321.
Report Division of Commercial Fishing for 1933-1934, 32-37.
- Cleveland, M. M.,** and **Fellers, C. R.:** Nutritive value of fish meals, laboratory and practical tests compared, 293-303.
- Cobb, Eben W.:** Report of librarian, 19.
- Culler, C. F.:** The future of Upper Mississippi fisheries, 328-331.
- Curtis, Brian:** Golden trout of Cottonwood lakes, 259-265.
- Davis, H. S.:** Chlorine gas effective in killing coarse fish (disc.), 280.
Growth and heredity in trout, 197-202.
Stream improvement, purpose and value, 63-67.
- Deason, Hilary J.,** joint author: see **Hile, Ralph.**
- Denmead, Talbott:** Suggested black bass legislation (1935), 97-102.
- Deuel, Charles R.:** Food cost of trout in New York State hatcheries, 172-175.
- Dinsmore, A. H.:** Age of trout in relation to time of spawning (disc.), 358-359.
Effect of heredity on growth of brook trout, 203-204.
Salmon run on the Skagit after experimental planting (disc.), 362.
- Embody, G. C.:** Trout eggs, relation of temperature to incubation period, 281-292.
- Embody, G. C.,** and **Sadler, W. O.:** Propagating daphnia and other forage organisms intensively in small ponds, 205-210.
- Faigenbaum, Harold M.:** Chemical characteristics of Adirondack lakes and ponds, 189-196.

- Faigenbaum**, Harold M., joint author: see **Moore**, Emmeline, et al.
- Faubion**, H. E., joint author: see **Wiebe**, A. H.
- Fellers**, C. R., joint author: see **Cleveland**, M. M.
- Fiedler**, R. H.: Codifying the fishery industry under the N. R. A., 32-36.
- Fish**, Frederic F.: Ulcer disease of trout, 252-258.
- Foerster**, R. E.: Fry production from eyed-egg planting, 379-381.
- Gordon**, Seth: Report of the secretary-treasurer for 1933-1934, 15-18.
- Greeley**, John R.: Heredity vs. environment as influence on time of spawning (disc.), 358.
- Greeley**, John R., joint author: see **Moore**, Emmeline, et al.
- Griffiths**, Francis P., and **Stansby**, Maurice E.: Significance of bacterial count and chemical tests in determining relative freshness of haddock, 401-406.
- Hachey**, H. B.: Weather predictions and coastal fisheries, 382-389.
- Hart**, John Lawson, and **Tester**, Albert L.: Herring spawning, quantitative studies, 307-312.
- Hayford**, Charles O.: Oxygen depletion in largemouth bass ponds caused by rapid decomposition of organic matter; control of vegetation in shallow lakes (disc.), 377.
- Hile**, Ralph, and **Deason**, Hilary J.: Growth of whitefish in Trout Lake, Wisconsin, 231-237.
- Hogan**, Joe: Largemouth black bass rearing at Lonoke, Arkansas, 127-131.
- Huntsman**, A. G.: Factors influencing return of salmon from the sea, 351-354.
- The problem of control in aquiculture, 364-367.
- James**, M. C.: Effect of 1934 drought on fish life, 57-62.
- Johnston**, Herbert: Can blood suckers affect the fish population of a lake? 363.
- Kingsbury**, Oliver R.: New bass hatchery at South Otselic, N. Y., and its first year's operations, 132-138.
- Kitson**, J. Arthur, joint author: see **Belding**, David L.
- Langlois**, T. H.: Environment vs. temperature as influence on time of spawning in rock bass (disc.), 359.
- New type of food grinder to prevent pulping of feed (disc.), 162.
- Social behavior of bass in rearing ponds, 146-150.
- Standard methods of computing bass production, 163-166.
- Locke**, S. B.: Value of clean streams, 87-92.
- Lord**, Russell F.: Hatchery trout as foragers and game fish, 339-345.
- Measuring the angler's catch (disc.), 52-53.
- MacKay**, H. H., and **Werner**, W. H. R.: Observations on maskinonge culture, 313-317.
- Markus**, Henry C.: Destruction of forage fish, 93-96.
- Growth of brook trout in hatcheries vs. growth when released (disc.), 80.
- M'Gonigle**, R. H.: Algae, a factor in some hatchery mortalities, 416-422.
- Meehan**, O. Lloyd: Fertilizers in pondfish production: ecological aspects, 151-154.
- Moore**, Emmeline, **Greeley**, J. R., **Greene**, C. W., **Faigenbaum**, H. M., **Nevin**, F. R., and **Townes**, H. K.: Problem in trout stream management, 68-80.
- Mottley**, C. McC.: Rainbow trout, origin and relation, 323-327.
- Needham**, P. R.: Quantitative studies of stream bottom foods, 238-247.
- Needham**, P. R., and **Taft**, A. C.: Observations on spawning of steelhead trout, 332-338.
- Nevin**, F. R.: joint author: see **Moore**, Emmeline, et al.
- Nobbs**, Percy E.: Measuring the angler's catch (disc.), 51-52.
- Odell**, T. T.: *Pomolobus pseudoharengus*, life history and ecology, 118-126.

- Parnell, I. W.:** Fish parasites and their importance, 390-399.
- Phillips, Mrs. J. T.:** Plankton distribution in Manitou and Brome lakes, Quebec, 461-463.
- Prevost, Gustave:** Fry and fingerling counting, new method and apparatus, 270-275.
- Potassium permanganate in fish culture, a criticism, 304-306.
- Radebaugh, Gus H., and Agersborg, H. P. Kjerschow:** Economic value of treated sewage effluent in wildlife conservation, 443-455.
- Reegan, C. C.:** Floating bass-brooding equipment, 143-445.
- Richardson, L. R.:** Mechanical as well as chemical aspects of flooding areas are to be considered (disc.), 367.
- Observations on the effects of dams on lakes and streams, 457-458.
- Richardson, L. R., and Belknap, G. W.:** A method of planting *Osmerus mordax* Mitchill on a small scale, 432-434.
- Roach, Lee S., and Wickliff, E. L.:** Relationship of aquatic plants to oxygen supply and their bearing on fish life, 370-376.
- Rodd, J. A.:** Report of the Committee on Foreign Relations, 20-23.
- Sadler, W. O.,** joint author: see **Emboddy, G. C.**
- Shillington, K. G.:** Selective breeding of speckled trout, 274-275.
- Smith, M. W.:** Dissolved oxygen content of fertilized waters, 408-415.
- Spier, J. D.:** Dominating species of aquatic plants in Brome and Manitou lakes, Quebec, 464-468.
- Stansby, Maurice E.,** joint author: see **Griffiths, Francis P.**
- Surber, Eugene:** Feeding of young bass in circular pools, 141.
- Surber, Thaddeus:** Measuring the angler's catch (disc.), 53.
- Taft, A. C.,** California steelhead experiments, 248-251.
- Taft, A. C.,** joint author: see **Needham, P. R.**
- Taylor, B. W., and Lindsay, R. C.:** Trout lakes in Gaspé County, Quebec, 424-431.
- Terrell, Clyde B.:** Copper sulphate to destroy coarse fish (disc.), 279-280.
- Tester, Albert L.,** joint author: see **Hart, John Lawson.**
- Tidd, Wilbur M.:** Gold fish and carp infested by *Lernaea carassii*, 176-180.
- Townes, H. K.,** joint author: see **Moore, Emmeline, et al.**
- Van Oosten, John:** Value of questionnaires in commercial fisheries regulations and surveys, 107-117.
- Webster, B. O.:** Report of the Vice-President of the Division of Fish Culture, 28-32.
- Werner, W. H. R.:** Artificial spawning of speckled trout, use of physiological saline, 346-349.
- Werner, W. H. R.,** joint author: see **MacKay, H. H.**
- White, H. C.:** Some facts and theories concerning Atlantic salmon, 360-362.
- Spawning period of brook trout, 356-357.
- Wickliff, E. L.:** Measuring the angler's catch (disc.), 53.
- Report of the American Fish Policy Committee, 25-27.
- State law regulating handling, transportation and selling of bait (disc.), 96.
- Survival of fish in water with oxygen content of .15 to 2.00 parts per million (disc.), 376-377.
- Wickliff, E. L.,** joint author: see **Roach, Lee S.**
- Wiebe, A. H.:** Nocturnal depressions in the dissolved oxygen in ponds, with special reference to excess of coarse vegetation and fertilizers, 181-188.
- Wiebe, A. H., Burr, J. C., and Faubion, H. E.:** Stream pollution in Texas with special reference to salt water from oil fields, 81-86.